

Automated Oxygen Control in Neonatology



Dr Peter Reynolds MB.BS MSc PhD

Consultant Neonatologist, St. Peter's Hospital, UK

Hon. Senior Lecturer Royal Holloway University of London

Joint Clinical Lead, Kent, Surrey & Sussex Neonatal Network

Acknowledgements and Disclosures

Thank you to the organisers for the invitation

I will discuss the neonatal use of automated oxygen control systems and have received financial support from the following companies

- Inspiration Healthcare plc (Oxygenie)
- Vapotherm (OAM)

I have received financial support from the following companies, but will not be discussing their products:

- Chiesi – Curosurf surfactant
- Biofloratech Ltd – Labinic probiotic

Background

SpO₂ is the best current continuous measure to achieve normoxia


NeOProM studies - 91-95% group had lower mortality than 85-89%


Hyperoxia increases the risk of acute and chronic lung disease, ROP, red blood cell oxidative stress, cardiac dysfunction and neurodevelopment abnormalities

Hypoxia increases NEC, cerebral palsy and mortality. Increased levels of damaging free radicals as oxygenation restored increases inflammation, and vascular remodelling occurs in the longer term

Intermittent Hypoxia


- No standard definition
- Alarm and averaging settings vary
- Staff perceptions of need for intervention vary
- Charts under-record event incidence (Brockmann PE 2013)






Damianos A, et al. Neonatal intermittent hypoxia persistently impairs lung vascular development and induces long-term lung mitochondrial DNA damage. J Appl Physiol. 2022 Nov 1;133(5):1031-1041

Frequency of IH events



Increase in IH events over time





mean+95% confidence interval


Important outcomes associated with IH

- Severe ROP (Di Fiore JM 2010)
- Sleep-disordered breathing (Di Fiore JM 2019)
- BPD (Fairchild KD 2019; Raffay TM 2019; Jensen EA 2021)
- Death or neurodevelopmental disability (Poets CF 2015)

BPD and IH – prediction of risk



Intermittent Hypoxia – Pre-Vent Study



(Ambalavanan et al 2023)

Control of Oxygen

- Control of oxygenation is part of our ventilation optimisation strategy
- It is not about just staying in the target range
- It is also about minimising hypoxic and hyperoxic episodes – frequency, depth and duration

LEONI PLUS
CLAC *inside*

Vapotherm®
Oxygen Assist Module


Evidence for effectiveness of O₂ Controllers



OxyGenie®
Inspired Oxygen Control

Sophie
SPQC *inside*

CLiO₂TM
AVEA®



(Dargaville PA 2021)

Automated vs manual oxygen control in preterm infants: SR & Meta-analysis

13 trials enrolling 343 preterm infants on respiratory support.

Auto controllers for the range SpO₂ 90-95%

- Increased time spent within the TR (p = .008)
- Reduced the time of hypoxia <80% (p<0.0001)
- Reduced the time of hyperoxia >98% (p = .009)
- Reduced number of manual adjustments (p = .002)



Attenuation of intermittent hypoxia events

Randomised cross-over study of automated oxygen control for preterm infants receiving nasal high flow

Peter R Reynolds,^{1,2} Thomas L Miller,^{3,4} Leonithas I Volakis,⁵ Nicky Holland,¹
George C Dungan,^{5,6} Charles Christoph Roehr,^{7,8} Kevin Ives⁷

- Prospective, two-centre, order-randomised, 24-hr crossover trial
- N=30 preterm infants requiring at least $\text{FiO}_2 \geq 25\%$ and at least 12 FiO_2 changes
- Vapotherm OAM ("Intello₂") algorithm

Category	Automated control	Manual control	P values*
Target SpO ₂ range [†]			
Episodes/hour (counts)	41 (32–52)	38 (32–46)	0.0387
Average episode duration (s)	70 (49–99)	48 (31–61)	<0.0001
Per cent time in range (%)	80 (70–87)	49 (40–57)	<0.0001
SpO ₂ >95% [‡]			
Episodes/hour (counts)	37 (30–46)	18 (13–25)	<0.0001
Average episode duration (s)	12 (9.5–14)	48 (34–68)	<0.0001
Per cent time in range (%)	12 (8.9–16)	23 (15–41)	<0.0001
SpO ₂ >98% [‡]			
Episodes/hour (counts)	4.5 (1.8–8.5)	5.5 (1.9–14)	0.572 (NS)
Average episode duration (s)	6.4 (5.1–7.0)	24 (20–32)	<0.0001
Per cent time in range (%)	0.68 (0.29–1.4)	3.7 (1.2–9.9)	<0.0001
SpO ₂ <range			
Episodes/hour (counts)	27 (22–30)	24 (21–26)	0.082 (NS)
Average episode duration (s)	17 (15–21)	42 (29–50)	<0.0001
Per cent time in range (%)	12 (8.8–17)	28 (17–36)	<0.0001
SpO ₂ <80% for ≥60 s			
Total number of episodes [§]	0 (0–1.3)	5 (2.8–14)	<0.0001
Average episode duration (s)	0 (0–69)	119 (97–148)	<0.0001





Clinical evaluation of a novel adaptive algorithm for automated control of oxygen therapy in preterm infants on non-invasive respiratory support

Gemma K Plottier,^{1,2} Kevin I Wheeler,³ Sanoj K M Ali,^{1,2} Omid Sadeghi Fathabadi,⁴ Rohan Jayakar,⁴ Timothy J Gale,⁴ Peter A Dargaville^{1,5}

- Prospective, single-centre, order-randomised, 24-hr crossover trial
- N=20 preterm infants requiring oxygen
- VDL 1.0 algorithm ("Oxygenie")

Table 1 Oxygen saturation (SpO₂) targeting

	Manual control	Automated control	p Value*
SpO ₂ in target range	55 (46–60)%	78 (75–87)%	0.0001
SpO ₂ below target range	19 (12–27)%	14 (7.8–19)%	0.0027
SpO ₂ above target range	25 (23–35)%	5.1 (3.1–6.9)%	0.0003
Eupoxia	56 (48–63)%	81 (76–90)%	<0.0001
SpO ₂ in alarm range (89%–96%)	81 (70–83)%	93 (90–98)%	0.0006
SpO ₂ in alarm range or higher when in air	81 (73–83)%	95 (92–98)%	<0.0001
SpO ₂ <80%	0.7 (0.10–1.3)%	0 (0–0.17)%	0.0006
SpO ₂ 80%–84%	2.6 (1.2–3.2)%	0.39 (0.10–0.67)%	0.0001
SpO ₂ 85%–88%	10 (6.8–15)%	3.5 (1.1–5.8)%	0.0002
SpO ₂ 97%–98% when in oxygen	5.0 (3.2–7.9)%	0.71 (0.28–1.5)%	0.0001
SpO ₂ 99%–100% when in oxygen	0.46 (0.22–1.4)%	0 (0–0.12)%	0.0010



Summary: Oxygenie vs OAM

	Hypoxia	Hyperoxia	In range
Manual	19-28%	23-25%	49-55%
OAM	12%	12%	80%
Oxygenie	14%	5.1%	78%

Author	Title	Journal
Kaltsogianni <i>et al</i>	Closed-loop oxygen system in late preterm/term, ventilated infants with different severities of respiratory disease	Acta Paediatrica 2023
Salverda <i>et al</i>	Clinical outcomes of preterm infants while using automated controllers during standard care (OxyGenie and CLiO2).	Arch Dis Child Fetal Neonatal Ed. 2023
Ali <i>et al</i>	Preliminary study of automated oxygen titration at birth for preterm infants.	Arch Dis Child Fetal Neonatal Ed. 2022
Salverda <i>et al</i>	Comparison of two devices for automated oxygen control in preterm infants: a randomised crossover trial.	Arch Dis Child Fetal Neonatal Ed. 2022
Dargaville <i>et al</i>	Automated control of oxygen titration in preterm infants on non-invasive respiratory support	Arch Dis Child Fetal Neonatal Ed. 2022
Salverda <i>et al</i>	The effect of automated oxygen control on clinical outcomes in preterm infants: a pre- and post-implementation cohort study	Eur J Pediatr. 2021
Sturrock S <i>et al</i>	A randomised crossover trial of closed loop automated oxygen control in preterm, ventilated infants	Acta Paediatrica. 2021
Plottier G <i>et al</i>	Clinical evaluation of a novel adaptive algorithm for automated control of oxygen therapy in preterm infants on non-invasive respiratory support	Arch Dis Child Fetal Neonatal Ed. 2017
Dargaville <i>et al</i>	Development and preclinical testing of an adaptive algorithm for automated control of inspired oxygen in the preterm infant	Arch Dis Child Fetal Neonatal Ed. 2017

Clinical Outcomes and AOC

FiO₂-C study

- Initial planned recruit 2340 infants <28 weeks gestation
- Randomisation to automated oxygen control with any CE-marked algorithm/device, or to routine manual oxygen control.

The primary outcomes are:

- death or any of severe ROP, BPD or NEC.
 - death or any of i) language/cognitive delay, ii) motor impairment, iii) severe visual impairment or iv) hearing impairment, at 2 years corrected gestational age.
-
- The trial commenced in July 2018; poor recruitment, has stopped early

CLiO₂TM vs OxygenieTM


	OXYGENIE	CLiO ₂	P
N=242	N=121	N=121	
Gestational Age	28+3	27+5	0.09
Birth Weight	1034	1011	0.60
Time in target range	71.5%	51.3%	<0.001
Hypoxia (SpO ₂ <80%) / Hyperoxia (SpO ₂ >98%)	0.7 / 1%	1.2 / 4%	<0.001
Invasive ventilation days, median (IQR)	0 (0–4.2)	2.1 (0–8.4)	0.012
CPAP days, median (IQR)	8.4 (4.8–19.8)	16.7 (6.3–31.1)	<0.001
Died, n (%)	7 (5.8)	3 (2.5)	0.34
Retinopathy of prematurity			
Early Treatment indicated n (%)	5 (4.5)	13 (11.2)	0.09
Received laser coagulation, n (%)	1 (0.9)	10 (8.6)	0.008
Intraventricular haemorrhage (≥stage 2), n (%)	27 (22.5)	18 (14.9)	0.21
Days in NICU, median (IQR)	28(15 – 42)	40(25 – 61)	<0.001

Cochrane Meta-analysis 2023

- 13 studies, 339 infants
- Improved time in target range
- Reduced hyperoxia
- Probably reduced hypoxia
- No clear effect on clinical outcomes

Frequency of hypoxic & hyperoxaemic episodes

- 58 babies <32 weeks studied on non-invasive ventilation
- Randomised to 24hrs on manual or automated O₂ control
- Episodes at extremes (>98% or <80%) were less common in automated control
- Long episodes (>30 seconds) were less common in automated control



Outcomes at 2-years

- Neurodevelopmental outcomes at 2 years of age were compared for infants born at 24–29 weeks gestational age before (2012–2015) and after (2015–2018)
- The primary outcome was a composite (death or severe impairment)
- 289 in the pre-AOC; 292 in the post-AOC epochs
- Outcome of mortality /severe NDI 17.9% pre-AOC (41/229) vs. 24.0% (47/196) post-AOC ($p = 0.12$)
- Limitations of retrospective study and use of different devices acknowledged

Why have trials not shown improved clinical outcomes?

- Outcomes depend on more than just oxygenation
- Influence of frequency and durations of IH events not tested in studies
- Rapid improvements in neonatal care influence outcomes
- Some adverse outcomes are rare, so very large sample sizes would be needed
- Risk of patients lost to follow-up in retrospective studies may not be random

Situations where auto-oxygen is helpful

- During handling: examinations, cares, nappy changes, parent skin-to-skin
- During procedures
 - Ultrasound scans
 - Umbilical / long line insertion
 - LISA procedure for surfactant
 - Cannulation / lumbar puncture / X-ray / Lung ultrasound etc
- Frequent desaturations (e.g. shunting) – compliance will still be better than manual control
- Where nursing staff are busy!

Summary

- Not all automated oxygen controllers are the same
- Characteristics "out of target range" should be considered
- Intermittent hypoxia and hyperoxia are (perhaps most) important
- Use of AOC should be throughout the "oxygen journey"
- Consider the additional benefits:
 - Reduction in nursing workload
 - Greater stability during procedures

Thank you

peter.reynolds1@nhs.net

X @neonataldoc

LinkedIn  dr-peter-reynolds